







STATO DELL'ARTE IN TERAPIA ONCOLOGICA: CARCINOMA POLMONARE

Lucio Buffoni


Oncologia Polmonare

AOU San Luigi Orbassano






Oncogene Addicted Tumor: Special Entity



ALGORITMO TERAPEUTICO NSCLC: 2017




Sc na

THE INTERN

Breakthrough of
Cancer
Immunot
T cells on the at

Antitumour immunity
enhanced by inhibiting
PD-L1/PD-1 and identifying
mutant neo-antigens

PAGES 496, 550, 563, 569, 572 & 577



Cancer is an immunological disorder

ORIGINAL ARTICLE

Nivolumab versus Docetaxel in Advanced Nonsquamous Non-Small-Cell Lung Cancer

H. Borghaei, L. Paz-Ares, L. Horn, D.R. Spigel, M. Soler, N.E. Ready, L.Q. Chow, E.E. Vokes, E. Felip, E. Holgado, F. Barlesi, M. Kohlhaufl, O. Arrieta, M.A. Burgio, J. Fayette, H. Lena, E. Poddighe, D.E. Gerber, S.N. Gettinger, C.M. Rudin, N. Rizvi, L. Crino, J. Brahmer, J. Estrella, J. Barlow, J. Carlezon, Jr., J. Hainsworth, J. Antonia, C. Dorange, C.T. Hainsworth, P. Graf Finckenstein, and J.R. Brahmer

ABSTRACT

BACKGROUND

Nivolumab, a fully human IgG4 programmed death 1 (PD-1) immune-checkpoint-inhibitor antibody, disrupts PD-1-mediated signaling and may restore antitumor immunity.

METHODS

In this randomized, open-label, international phase 3 study, we assigned patients with nonsquamous non-small-cell lung cancer (NSCLC) that had progressed during or after platinum-based doublet chemotherapy to receive nivolumab at a dose of 3 mg per kilogram of body weight every 2 weeks or docetaxel at a dose of 75 mg per square meter of body-surface area every 3 weeks. The primary end point was overall survival.

PEER REVIEW

ACCEPT YOUR
OWN PAPER
How some scientists are
duping the system
PAGE 400


MICROSCOPY

THE CA
AIMING
Atomic resolu
for the
PAGE

Check-Mate 17

Nivolumab versus Docetaxel in Advanced Squamous-Cell Non-Small-Cell Lung Cancer

OS



No. at Risk


Nivolumab	135	113	86	69	52	31	15	7	0
Docetaxel	137	103	68	45	30	14	7	2	0

Check-Mate 57

Nivolumab versus Docetaxel in Advanced Nonsquamous Non-Small-Cell Lung Cancer

OS

Overall Survival



No. at Risk


Nivolumab	292	232	194	169	146	123	62	32	9	0
Docetaxel	290	244	194	150	111	88	34	10	5	0



Key-Note 10

Pembrolizumab versus docetaxel for previously treated, PD-L1-positive, advanced non-small-cell lung cancer (KEYNOTE-010): a randomised controlled trial


ALL PTS



	0	5	10	15	20	25
Number at risk						
Pembrolizumab 2 mg/kg	344	259	115	49	12	0
Pembrolizumab 10 mg/kg	346	255	124	56	6	0
Docetaxel	343	212	79	33	1	0

OS

PD-L1
≥50%



	0	5	10	15	20	25
Number at risk						
Pembrolizumab 2 mg/kg	139	110	51	20	3	0
Pembrolizumab 10 mg/kg	151	115	60	25	1	0
Docetaxel	152	90	38	19	1	0



ALGORITMO TERAPEUTICO NSCLC: 2017

NSCLC IIB-IV

EGFR e ALK WT

PS 0-1/Età < 70

Non-SCC

SCC

CT + **Beva**
P - Pem

P - Gem
P - Doc

Docetaxel +/-
Nintedanib


Docetaxel

Nivolumab
Pembrolizumab PDL1>1%

2016-17



KEYNOTE 024: STUDY DESIGN



Key End Points

Primary: PFS (RECIST v1.1 per blinded, independent central review)

Secondary: OS, ORR, safety

Exploratory: DOR

Platinum-Doublet Chemotherapy Options


- Pemetrexed 500 mg/m² on day 1 of 21-day cycles + Carboplatin AUC 5 or 6 on day 1 of 21-day cycles
 - Pemetrexed 500 mg/m² on day 1 of 21-day cycles + Cisplatin 75 mg/m² on day 1 of 21-day cycles
 - Paclitaxel 200 mg/m² on day 1 of 21-day cycles + Carboplatin AUC 5 or 6 on day 1 of 21-day cycles
 - Gemcitabine 1250 mg/m² on days 1 and 8 of 21-day cycles + Carboplatin AUC 5 or 6 on day 1 of 21-day cycles
 - Gemcitabine 1250 mg/m² on days 1 and 8 of 21-day cycles + Cisplatin 75 mg/m² on day 1 of 21-day cycles
- Nonsquamous NSCLC only

Reck M, ESMO 2016




I LINE: SINGLE IMMUNO AGENT VS STANDARD CT

Overall Survival



Progression Free Survival



Reck M, NEJM 2016; Reck M, ESMO 2016



ALGORITMO TERAPEUTICO NSCLC: 2017

NSCLC IIB-IV

EGFR e ALK WT

EGFR e ALK WT

PDL1 > 50%

PS 0-1/Età < 70

~~CHEMIOTERAPIA~~

Non-SCC

SCC

CT + **Beva**
P - Pem

P - Gem
P - Doc

Docetaxel +/-
Nintedanib

Docetaxel


Nivolumab
Pembrolizumab PDL1 > 1%

PEMBROLIZUMAB

CHEMIOTERAPIA



CHECKMATE 057: RESPONSE AND DURATION OF RESPONSE




Variable	Nivolumab (N = 292)	Docetaxel (N = 290)
Objective response†		
No. of patients	56	36
% of patients (95% CI)	19 (15–24)	12 (9–17)
Estimated odds ratio (95% CI)	1.7 (1.1–2.6)	
P value	0.02	
Best overall response — no. (%)		
Complete response	4 (1)	1 (<1)
Partial response	52 (18)	35 (12)
Stable disease	74 (25)	122 (42)
Progressive disease	129 (44)	85 (29)
Could not be determined	33 (11)	47 (16)
Time to response — mo‡§		
Median	2.1	2.6
Range	1.2–8.6	1.4–6.3
Duration of response — mo‡¶		
Median	17.2	5.6
Range	1.8 to 22.6+	1.2+ to 15.2+


Borghaei H, NEJM 2015



Supportive PRO Analysis: Change From Baseline to Week 15 EORTC QLQ-C30 Functioning and Symptom Scales



IMMUNE-RELATED AEs WITH IMMUNOTHERAPY



ALGORITMO TERAPEUTICO NSCLC: I LINEA 2017

NSCLC IIB-IV

CHEMIO**2008**TERAPIA

55%

EGFR WT
ALK NT
ROS1 NT
Età media più alta
All histologies
Fumatori/Ex fumatori

IMMUNO**2017**TERAPIA

25%

EGFR WT
ALK/ROS1 NT
PDL1 > 50%
Età media più alta
All histologies
Fumatori

TARGET**2009**TERAPIES

20%

EGFR mut+
ALK traslocato
ROS1+
Età media più bassa
Adenocarcinoma
Non fumatori



CONCLUSIONI NSCLC 2017...

- **Non Oncogene Addicted:**

Immunotherapy is a standard of care with IMPRESSIVE efficacy, rapid ONSET, LONG effect, MILD and PECULIAR toxicities

- **FUTURE:**

- PREDICTIVE BIOMARKERS (PDL1 + TMB)

- COMBINATIONS with CHEMOTHERAPY or IMMUNOTHERAPY (Anti PDL1/Anti CTLA4)

-

- **Oncogene Addicted:**

EGFR +:

- TKIs I gen -> TKIs 3 gen PFS 24 mesi e OS 3-4 anni

- TKIs 3 gen in I line to prevent PD and Brain Mets?

- **ALK +:**

- Crizo I line -> Alectinib/Ceritinib -> Lorlatinib/Brigatinib PFS 20 mesi OS 4 anni

- Alectinib I line to prevent PD and Brain Mets?

